arXiv:0801.2075v1 [math.DG] 14 Jan 2008

NEUTRAL BI-HERMITIAN GRAY SURFACES.

Wlodzimierz Jelonek

Abstract. The aim of this paper is to give examples of compact neutral 4-manifolds (M,g) whose Ricci tensor ρ satisfies the relation $\nabla_X\rho(X,X)=\frac{1}{3}X\tau g(X,X)$. We present also a family of new Einstein bi-Hermitian neutral metrics on ruled surfaces of genus $g>1$.

This paper I dedicate to Professor Kouei Sekigawa on his sixtieth birthday.

0. Introduction. In the present paper we are concerned with the class of neutral semi-Riemannian 4-manifolds (M,g) whose Ricci tensor ρ satisfies the condition

$$(*)\qquad \nabla_X\rho(X,X)=\frac{1}{3}X\tau g(X,X)$$

where τ is the scalar curvature of (M,g). These class of manifolds was introduced by A. Gray (see [G],[Be]). For general facts and some results concerning neutral 4-manifolds we refer to [K], [Pe],[A], [D],[M-L]. In [M] the condition for the existence of two opposite almost complex structures on 4-manifolds are studied. Note that in [A] (p.187) it is stated that there are no known neutral Hermitian-Einstein metrics on ruled surfaces except trivial ones. In this note we give among others many new, non-trivial examples of such manifolds. The main subject of the paper is the construction of new Einstein and Gray neutral metrics on ruled surfaces. The methods however are very similar to those used by me in [J-1]-[J-4].

In our papers [J-2]-[J-4] we have described Riemannian metrics g on compact complex surfaces (M,J) such that (M,g) satisfies the condition $*$ and has J-invariant Ricci tensor. In particular we have given a complete classification of bi-Hermitian Gray surfaces on ruled surfaces of genus $g\geq 1$ and surfaces which are of co-homogeneity 1 on ruled surfaces of genus $g=0$. We denote by g also the Riemannian metric but it should not cause any misunderstandings. The technics

---

MS classification:53C05,53C20,53C25. The paper was partially supported by KBN grant 2P0 3A 02324.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

used in [J-3] can be applied to describe a large class of neutral bi-Hermitian Gray surfaces. The equations, describing such surfaces whose Ricci tensor has exactly two real eigenvalues such that the corresponding two dimensional eigendistributions are space-like and time-like, are described by the same equations as in [J-3],[J-4]. In that way we present a large class of explicit examples of neutral bi-Hermitian Gray surfaces. In particular we get a large class of Einstein neutral bi-Hermitian surfaces on ruled surfaces of genus $g > 1$.

1. Neutral $\mathcal{A}C^{\perp}$-surfaces. By an $\mathcal{A}C^{\perp}$- manifold (see [Be],[G]) we mean a semi-Riemannian manifold (M, g) satisfying the condition

$$(*) \qquad \mathfrak{C}_{XYZ}\nabla_X\rho(Y,Z) = \frac{2}{(\dim M + 2)}\mathfrak{C}_{XYZ}X\tau g(Y,Z),$$

where ρ and τ are the Ricci tensor and the scalar curvature of (M, g) respectively and $\mathfrak{C}$ means the cyclic sum. In this paper, an $\mathcal{A}C^{\perp}$-manifold with neutral metric is also called a neutral Gray manifold. A Riemannian manifold (M, g) is an $\mathcal{A}C^{\perp}$ manifold if and only if the Ricci endomorphism Ric of (M, g) is of the form $Ric = S + \frac{2}{n+2}\tau Id$ where S is a Killing tensor and $n = \dim M$. Let us recall that a symmetric (1,1) tensor S on a semi-Riemannian manifold (M, g) is called a Killing tensor if $g(\nabla S(X,X), X) = 0$ for all $X \in TM$ and that a semi-Riemannian manifold whose Ricci tensor is a Killing tensor is called an $\mathcal{A}$-manifold.

On a semi-Riemannian manifold (M, g) a distribution $\mathcal{D} \subset TM$ is called umbilical (see [J-3]) if $\mathcal{D}$ is non-degenerate (i.e. a metric g is non-degenerate on $\mathcal{D}$) and $\nabla_X X_{|\mathcal{D}^{\perp}} = g(X,X)\xi$ for every $X \in \Gamma(\mathcal{D})$, where $X_{|\mathcal{D}^{\perp}}$ is the $\mathcal{D}^{\perp}$ component of X with respect to the orthogonal decomposition $TM = \mathcal{D} \oplus \mathcal{D}^{\perp}$. The vector field ξ is called the mean curvature normal of $\mathcal{D}$. The foliation tangent to involutive distribution $\mathcal{D}$ is called totally geodesic if its every leaf is a totally geodesic (i.e $\nabla_X X \in \mathcal{D}$ for any $X \in \mathcal{D}$) non-degenerate submanifold of (M, g). In the sequel we shall not distinguish between $\mathcal{D}$ and a foliation tangent to $\mathcal{D}$ and we shall also say that $\mathcal{D}$ is totally geodesic in such a case.

It is not difficult to prove exactly as in the Riemannian case (see [J-3]) the following lemma:

Lemma 1. *Let $S \in End(TM)$ be a (1,1) tensor on a neutral semi-Riemannian 4-manifold (M, g). Let us assume that S has exactly two everywhere different eigenvalues λ, μ of the same multiplicity 2, i.e. dim $\mathcal{D}_\lambda$= dim $\mathcal{D}_\mu = 2$, where $\mathcal{D}_\lambda, \mathcal{D}_\mu$ are non-degenerate eigendistributions of S corresponding to λ, μ respectively. Then S is a Killing tensor if and only if both distributions $\mathcal{D}_\lambda$ and $\mathcal{D}_\mu$ are umbilical with mean curvature normal equal respectively*

$$\xi_\mu = \frac{\nabla\mu}{2(\lambda-\mu)},\ \xi_\lambda = \frac{\nabla\lambda}{2(\mu-\lambda)}.$$

We shall call a bi-Hermitian surface with bi-Hermitian Ricci tensor simply as bi-Hermitian surface If (M, g) is also an $\mathcal{A}C^{\perp}$ manifold, then we call it a neutral bi-Hermitian Gray surface.

Proposition 1. *Let us assume that (M, g) is a simply connected neutral Gray 4-manifold and the Ricci tensor S ($\rho(X,Y) = g(SX,Y)$) has exactly two real* *eigenvalues* λ, μ *and no null eigenvectors. Then there exist two Hermitian complex structures* $J, \bar{J}$ *commuting with* S *and* (M, g) *is a bi-Hermitian neutral Gray surface.*

Proof. Analogous to [J-1].♢
We shall prove

Proposition 2. *Let* (M, g) *be a 4-dimensional neutral semi-Riemannian manifold. Let* $\mathcal{D}$ *be a two dimensional totally geodesic bundle-like and space-like foliation on* M*. Then* M *admits (up to 4-fold covering) two opposite Hermitian structures* $J, \bar{J}$ *such that* $J|\mathcal{D} = -\bar{J}|\mathcal{D}, J|\mathcal{D}^{\perp} = \bar{J}|\mathcal{D}^{\perp}$*. The nullity distributions of both* $J, \bar{J}$ *contain* $\mathcal{D}$*.*

Proof. To prove the first part of the Proposition it is enough to show that on M there exists a Killing tensor with eigendistributions $\mathcal{D}, \mathcal{D}^{\perp}$. Since the foliation $\mathcal{D}$ is totally geodesic and bundle-like it follows that $\nabla_X X \in \Gamma(\mathcal{D})$ (resp.) $\nabla_X X \in \Gamma(\mathcal{D}^{\perp})$ if $X \in \Gamma(\mathcal{D})$ (resp.) $X \in \Gamma(\mathcal{D}^{\perp})$. Consequently a tensor S defined by

$$SX = \lambda X \text{ if } X \in \mathcal{D}$$
$$SX = \mu X \text{ if } X \in \mathcal{D}^{\perp}$$

where $\lambda \neq \mu$ are two different real numbers is a smooth Killing tensor. We can assume (up to 4-fold covering) that the distributions $\mathcal{D}, \mathcal{D}^{\perp}$ are orientable. Let us denote by J the only almost Hermitian structure which preserves $\mathcal{D}, \mathcal{D}^{\perp}$ and agrees with their orientations. We define $\bar{J}$ by $J|\mathcal{D} = -\bar{J}|\mathcal{D}, J|\mathcal{D}^{\perp} = \bar{J}|\mathcal{D}^{\perp}$. From Proposition 1 it follows that both $J, \bar{J}$ are Hermitian. Now let $\{E_1, E_2\}$ be a local orthonormal frame on $\mathcal{D}$. Then

$$\nabla J(E_1, E_1) + J(\nabla_{E_1} E_1) = \nabla_{E_1} E_2.$$

It follows that $\nabla J(E_1, E_1) \in \mathcal{D}$ and consequently $\nabla_{E_1} J = 0$. Analogously $\nabla_{E_2} J = 0$. ♢

Now we give a theorem, whose proof is analogous as in the Riemannian case (see [M-S],[B],[S]).

Theorem 1. *Let us consider the manifold* $U = (a, b) \times P$*, where* (P, g_0) *is a 3-dimensional semi-Riemannian* $\mathcal{A}$*-manifold (a principle* S^1 *bundle* $p : P \to \Sigma$ *) over a Riemannian surface* (Σ, g_{can}) *of constant sectional curvature* K*, with an* $\mathcal{A}C^{\perp}$*-metric*

$$g = dt^2 + f(t)^2\theta^2 - g(t)^2 p^* g_{can}, \tag{1.5}$$

where $g_0 = \theta^2 - p^* g_{can}$*,* $f, g \in C^{\infty}(a, b)$ *and* θ *is the connection form of* P*. Then the metric* g *on* U *extends to a smooth* $\mathcal{A}C^{\perp}$ *metric on the ruled surface* M *which is a* $\mathbb{C}P^1$*-bundle over* Σ *and such that* U *is an open and dense subset of* M *if and only if the functions* $f, g \in C^{\infty}(a, b)$ *satisfy the conditions:*
(a) $f(a) = f(b) = 0, f'(a) = 1, f'(b) = -1,$
(b) $g(a) \neq 0 \neq g(b), g'(a) = g'(b) = 0.$

Remark. Let us note that the metric (1.5) induces a semi-Riemannian metric on M if the functions f, g satisfy:

(i) f is positive on (a,b) , and f is odd at a and b , i.e. f is the restriction of a function f on $\mathbb{R}$ satisfying $f(a+t)=-f(a-t), f(b+t)=-f(b-t)$;

(ii) g is positive on $[a,b]$ and even at a and b which means that g is the restriction of a function g such that $g(a+t)=g(a-t), g(b+t)=g(b-t)$;

The proof is similar to the description of the metric in polar coordinates (see [Be] Lemma 9.114 and Theorem 9.125.)

Note that it is an easy exercise to prove that functions f,g satisfying the ODE characterizing $\mathcal{A}C^{\perp}$-metrics together with the initial conditions $(a),(b)$ extend to respectively odd and even (with respect to the points a,b) real analytic functions around a,b. The fact that the conditions $(a),(b)$ of Theorem 1 imply (i) and (ii) is based on the following elementary Lemma.

Lemma 2. *Let us assume that the function Q is real analytic in a closed interval $[x_0,x_1]$, Q is positive in (x_0,x_1) and $Q(x_0)=Q(x_1)=0, Q'(x_0)\neq 0, Q'(x_1)\neq 0$. Let a real function $\phi(t)$ satisfy an equation*

$$\phi''=\frac{1}{2}Q'(\phi), \phi(0)=x_1, \phi'(0)=0, \tag{1.6}$$

Then $(\phi')^2=Q(\phi)$ and ϕ is a real analytic and periodic function $\phi\in C^{\omega}(\mathbb{R})$ satisfying conditions $\phi(0)=x_0, \phi(l)=x_1$ where l is the first positive real number such that $\phi(l)=x_1$. Also ϕ is an even function at $t=0$ and $t=l$.

Proof. The first part of Lemma is well known (see [B],p.445, Lemma 16.37). If ϕ satisfies equation (1.6) then $(\phi')^2=Q_0(\phi)$ where Q_0 is a primitive function of Q'. From the boundary conditions $Q_0=Q$. It is enough to show that ϕ is even at $t=0$ (for $t=l$ the proof is similar) which is equivalent to $\phi^{(2k-1)}(0)=0$ for every $k\in\mathbb{N}$. We prove by induction with respect to k that for every $F\in C^{\omega}(\mathbb{R})$ the function $\psi(t)=F(\phi(t))$ is even at $t=0$. If $k=1$ then $\psi'(0)=F'(\phi(0))\phi'(0)=0$. Let us assume that the result holds for $2k-1$. Let us write $\frac{1}{2}Q'=Q_1$. We get

$$\begin{aligned}\psi^{(2k+1)}&=(\psi'')^{(2k-1)}=(F''(\phi)(\phi')^2+F'(\phi)\phi'')^{(2k-1)}\\&=(F''(\phi)Q(\phi)+F'(\phi)Q_1(\phi))^{(2k-1)}=\end{aligned}$$
$$\sum\frac{(2k-1)!}{p!(2k-1-p)!}((F''(\phi))^{(p)}Q(\phi)^{(2k-1-p)}+(F'(\phi))^{(p)}(Q_1(\phi))^{(2k-1-p)}).$$

Thus $\psi^{(2k+1}(0)=0$. It follows that $F(\phi)$ is even at 0 for every F. If $F=id$ then we get that ϕ is even at $0.\diamondsuit$

Using this lemma one can show that in every case considered below in the paper function g is even and f is odd. Both functions are non-negative and g is positive which follows from the formula on g.

2. Neutral bi-Hermitian Gray surfaces. In this section we shall construct bi-Hermitian neutral metrics on ruled surfaces $M_{k,g}$ of genus g. The ruled surface $(M_{k,g},g)$ is locally of co-homogeneity 1 with respect to the group of all local isometries of $(M_{k,g},g)$ and an open, dense submanifold $(U_{k,g},g)\subset(M_{k,g},g)$ is isometric to the manifold $(a,b)\times P_k$ where (P_k,g_k) is a 3-dimensional semi-Riemannian $\mathcal{A}$-manifold (a principal circle bundle $p:P_k\rightarrow\Sigma_g$) over a Riemannian surface (Σ_g,g_{can}) of constant sectional curvature $K\in\{-4,0,4\}$ with a metric

$$g_{f,g}=dt^2+f(t)^2\theta^2-g(t)^2p^*g_{can}, \tag{2.1}$$

where $g_k = \theta^2 - p^* g_{can}$ and θ is the connection form of P_k such that $d\theta = 2\pi k\, p^*\omega$, where the de Rham cohomology class $[\omega] \in H^2(\Sigma_g, \mathbb{R})$ defined by the form ω is an integral class corresponding to the class $1 \in H^2(\Sigma_g, \mathbb{Z}) = \mathbb{Z}$. Let $\theta^\sharp$ be a vector field dual to θ with respect to g_k. Let us consider a local orthonormal frame $\{X, Y\}$ on (Σ_g, g_{can}) and let X^h, Y^h be horizontal lifts of X, Y with respect to $p : M_{k,g} \to \Sigma_g$ (i.e. $dt(X^h) = \theta(X^h) = 0$ and $p(X^h) = X$) and let $H = \frac{\partial}{\partial t}$. Let us define two almost Hermitian structures $J, \bar{J}$ on M as follows

$$JH = \frac{1}{f}\theta^\sharp, JX^h = Y^h,\ \bar{J}H = -\frac{1}{f}\theta^\sharp, \bar{J}X^h = Y^h.$$

Proposition 3. *Let $\mathcal{D}$ be a distribution spanned by the fields $\{\theta^\sharp, H\}$. Then $\mathcal{D}$ is a non-degenerate totally geodesic foliation with respect to the metric $g_{f,g}$ where g is a non constant function. Both structures $J, \bar{J}$ are Hermitian and $\mathcal{D}$ is contained in the nullity of J and $\bar{J}$. The distribution $\mathcal{D}^\perp$ is umbilical with the mean curvature normal $\xi = -\nabla \ln g$. Let λ, μ be eigenvalues of the Ricci tensor S of $g_{f,g}$ corresponding to eigendistributions $\mathcal{D}, \mathcal{D}^\perp$ respectively. Then the following conditions are equivalent:*

(a) There exists $D \in \mathbb{R}$ such that $\lambda - \mu = Dg^2$,
(b) There exist $C, D \in \mathbb{R}$ such that $\mu = Dg^2 - C$,
(c) $\lambda - 2\mu$ is constant,
(d) $(U_{k,g}, g_{f,g})$ is a neutral bi-Hermitian Gray surface.

Proof. The first assertion of Proposition 3 is a consequence similar to Proposition 3 in [J-2]. Note that $\nabla\lambda = H\lambda H, \nabla\mu = H\mu H$. Consequently $tr_g \nabla S = \frac{1}{2}\nabla\tau = (H\lambda + H\mu)H$. On the other hand one can easily check that $tr_g \nabla S = 2(\mu - \lambda)\xi + H\lambda H$. Thus

$$\nabla\mu = 2(\lambda - \mu)\nabla \ln g.$$

Now we prove that (a) $\Rightarrow$ (b). If (a) holds then $\nabla\mu = 2Dg^2\frac{\nabla g}{g} = D\nabla g^2$. Thus $\nabla(\mu - Dg^2) = 0$ which implies (b).

(b)$\Rightarrow$ (a). We have

$$-\frac{\nabla g}{g}2(\mu - \lambda) = \nabla\mu = 2Dg\nabla g,$$

and consequently $\nabla g(\frac{(\mu-\lambda)}{g} - Dg) = 0$ which is equivalent to (a).

(a)$\Rightarrow$(c). We have $\lambda - \mu = Dg^2$ and consequently $\nabla\mu = 2Dg\nabla g = D\nabla g^2$. Thus $\nabla\lambda = \nabla(\mu + Dg^2) = 2D\nabla g^2$ and $\nabla\lambda - 2\nabla\mu = 0$ which gives (c).

(c)$\Rightarrow$ (a). If $\nabla\lambda = 2\nabla\mu$ then $\nabla\lambda = 4(\lambda - \mu)\frac{\nabla g}{g}$. Consequently $\nabla\lambda - \nabla\mu = 2(\lambda - \mu)\frac{\nabla g}{g}$ and $\nabla \ln|\lambda - \mu| = 2\frac{\nabla g}{g} = 2\nabla \ln g$, which means that $\nabla \ln|\lambda - \mu| g^{-2} = 0$ or $\lambda = \mu$ on the whole of M. In the last case we obtain an Einstein metric. It follows that $\ln\frac{|\lambda-\mu|}{g^2} = \ln D$ for some $D \in \mathbb{R}_+$ or $\lambda - \mu = 0$ (D=0), which is equivalent to (a).

(d)$\Leftrightarrow$(c). This equivalence follows from [J-3]. $\diamondsuit$

Theorem 2. *On any ruled surface $M_{k,g}$ of genus g with $k > 0$ there exist a one-parameter family of neutral bi-Hermitian $\mathcal{AC}^\perp$-metrics $\{g_x : x \in (0, \epsilon_s)\}$,* *where* $\epsilon_s > 0$ *depends only on* g *and* k*, which consists of neutral bi-Hermitian Gray metrics on* $M_{k,g}$*.*

Proof. Note that for the first Chern class $c_1(\Sigma_g) \in H^2(\Sigma_g, \mathbb{Z})$ of the complex curve Σ_g we have the relation $c_1(\Sigma_g) = \chi\alpha$, where $\alpha \in H^2(\Sigma_g, \mathbb{Z})$ is an indivisible integral class and $\chi = 2 - 2g$ is the Euler characteristic of Σ_g. Let us write $s = \frac{2k}{|\chi|}$ if $g \neq 1$ and $s = k$ if $g = 1$. Then it is easy to show using O'Neill formulas for a semi-Riemannian submersion (see [ON],[B],[S]) that the manifold $(M_{k,g}, g)$ with the metric g given by (2.1) has the Ricci tensor with the following eigenvalues :

$$\lambda_0 = -2\frac{g''}{g} - \frac{f''}{f}, \tag{2.2a}$$

$$\lambda_1 = -\frac{f''}{f} - 2\frac{f'g'}{fg} + 2s^2\frac{f^2}{g^4}, \tag{2.2b}$$

$$\lambda_2 = -\frac{g''}{g} - \frac{f'g'}{fg} - (\frac{g'}{g})^2 - 2s^2\frac{f^2}{g^4} - \frac{K}{g^2}, \tag{2.2c}$$

where λ_0, λ_1, correspond to eigenfields $T = \frac{d}{dt}, \theta^\sharp$ and λ_2 corresponds to a two-dimensional eigendistribution orthogonal to T and $\theta^\sharp$. Note that the equations we have got are practically the same as equations for Gray manifolds obtained in the Riemannian case. The only difference is that we now have $-K$ instead of K, so the cases $K > 0$ and $K < 0$ are now reversed, which give the different geometric meaning to the equations in Riemannian and semi-Riemannian neutral cases. We present here the method of solving equations (2.2) for the convenience of the reader and completeness, although the calculations are just the same as in [J-4]. If $(M, g) \in \mathcal{AC}^\perp$ is a bi-Hermitian Gray surface then $\lambda_0 = \lambda_1 = \lambda$, and if we denote $\mu = \lambda_2$, Proposition 3 b implies an equation

$$\mu = Dg^2 - C \tag{2.3}$$

for some $D, C \in \mathbb{R}$. Since $\lambda_0 = \lambda_1$ we get

$$f = \pm\frac{gg'}{\sqrt{s^2 + Ag^2}}. \tag{2.4}$$

Using a homothety of the metric we can assume that $A \in \{-1, 0, 1\}$. In the case $A = 0$ we get a neutral Kähler metric and these metrics on compact complex surfaces we shall describe in section 4. So we restrict our considerations to the case $A \in \{-1, 1\}$. Now we introduce a function h such that $h^2 = s^2 + Ag^2$. Note that $\mathrm{im}h \subset (-s, s)$ if $A = -1$, and $\mathrm{im}h \subset (s, \infty)$ if $A = 1$. Then $g = \sqrt{|s^2 - h^2|}$. Let us introduce a function z such that $h' = \sqrt{z(h)}$. Note that

$$f = h' \text{ and } f' = \frac{1}{2}z'(h). \tag{2.5}$$

It follows that equation (2.3) is equivalent to

$$z'(h) - z(h)\frac{s^2 + h^2}{h(s^2 - h^2)} = \frac{4\epsilon}{h} + \frac{D(s^2 - h^2)^2}{h} - \frac{C(s^2 - h^2)}{h}, \tag{2.6}$$

where $\epsilon = -\mathrm{sgn} KA \in \{-1, 0, 1\}$. It follows that

$$z(h) = (1-(\frac{h}{s})^2)^{-1}(-4\epsilon(\frac{h}{s})^2 - \frac{Ds^4}{5}(\frac{h}{s})^6 + (Ds^4 - \frac{Cs^2}{3})(\frac{h}{s})^4 + \\ +(2Cs^2 - 3Ds^4)(\frac{h}{s})^2 - 4\epsilon + Cs^2 - Ds^4 + \frac{E}{s}\frac{h}{s}). \tag{2.7}$$

Let us denote again $C = Cs^2, D = Ds^4, E = \frac{E}{s}$ and let

$$z_0(t) = (1-t^2)^{-1}(-4\epsilon(1+t^2) + D(-\frac{1}{5}t^6 + t^4 - 3t^2 - 1) + C(-\frac{1}{3}t^4 + 2t^2 + 1) + Et). \tag{2.8}$$

Write

$$P(t) = -4\epsilon t^2 - \frac{D}{5}t^6 + (D - \frac{C}{3})t^4 + (2C - 3D)t^2 + Et - 4\epsilon + C - D. \tag{2.9}$$

Then $z_0(t) = \frac{P(t)}{1-t^2}$. Note that $z(h) = z_0(\frac{h}{s})$ and $z'(h) = \frac{1}{s}z_0'(\frac{h}{s})$. In view of Th. 1. we are looking for real numbers $x, y \in \mathbb{R}, x > y$ such that

$$z_0(x) = 0, z_0'(x) = -2s, \tag{2.10a}$$

$$z_0(y) = 0, z_0'(y) = 2s, \tag{2.10b}$$

and $z_0(t) > 0$ for $t \in (y, x)$. Note that equations (2.10a) are equivalent to

$$-4\epsilon x^2 - \frac{D}{5}x^6 + (D - \frac{C}{3})x^4 + (2C - 3D)x^2 - 4\epsilon + C - D + Ex = 0 \tag{2.11a}$$

$$-8\epsilon x - \frac{6D}{5}x^5 + 4(D - \frac{C}{3})x^3 + 2(2C - 3D)x + E = -2s(1 - x^2). \tag{2.11b}$$

Equations (2.11) yield

$$D = \frac{5(-3E - 6s - 24\epsilon x + 3Ex^2 - 12sx^2 - 8\epsilon x^3 + 2sx^4)}{2(-1+x)x(1+x)(15 + 10x^2 - x^4)}, \tag{2.12a}$$

(2.12b)

$$C = \frac{3(5E + 10s + 80\epsilon x + 30sx^2 - 10Ex^2 + 5Ex^4 - 10sx^4 - 16\epsilon x^5 + 2sx^6)}{2(-1+x)x(1+x)(-15 - 10x^2 + x^4)}$$

Solving in a similar way equations (2.10b) one can see that there exists a function z_0 satisfying the equations (2.10) if

$$(x+y)(-4\epsilon(-5x + x^3 + 5y + 2x^2y - 2xy^2 - y^3) \\ +s(5 + 2x^3y + 2xy^3 + 3y^2 + 3x^2 + x^2y^2 - 16xy)) = 0, \tag{2.13}$$

where $x > y$, $x, y \in (-1, 1)$ in the case $A = -1$ and $x, y \in (1, \infty)$ in the case $A = 1$. Using standard methods one can check that in the case of the genus $g \leq 1$ (i.e. if $K = 4$ or $K = 0$) the only solutions of (2.13) giving a positive function z are $x = -y \in (0, 1)$. In the case of genus $g \geq 2$ we have $K = -4$ and apart from the solutions with $x = -y$ (see [J-3]) there is an additional family of solutions with $\epsilon = -1$ for

$s \in (0,\eta)$ on $M_{k,g}$ where $\eta = \frac{2}{17}(15+4\sqrt{13})\sqrt{\frac{1}{3}(13(10\sqrt{13}-35)))} \simeq 2.05318...$ One can check that there exist families of solutions with $-1<y<x<1$ for $k \in \mathbb{N}$ such that $k<(g-1)\eta$. In fact it is not difficult to check that if $g>1$ then on the sets $F=\{(x,y): -1<y<x<1\}$ and $H=\{(x,y): 1<y<x\}$ the function $G(x,y) = -4\epsilon(-5x+x^3+5y+2x^2y-2xy^2-y^3)+s(5+2x^3y+2xy^3+3y^2+3x^2+x^2y^2-16xy)$ is negative and positive somewhere on the boundaries of both of F and H for $s<2$. In the case $g \leq 1$ G is positive on both F and H. We have

$$\begin{aligned} G_x(x,y) &= -4\epsilon(-5+3x^2+4xy-2y^2)+2s(3x^2y+y^2x+3x-8y+y^3),\\ G_y(x,y) &= -4\epsilon(5+2x^2-4xy-3y^2)+2s(x^3+3y+x^2y-8x+3x^2y). \end{aligned}$$

The equations $G_x=0, G_y=0$ are equivalent to

$$\begin{gathered} -4\epsilon(-5+3x^2+4xy-2y^2)+2s(3x^2y+y^2x+3x-8y+y^3)=0,\\ (x+y)(10\epsilon(y-x)+s(-5+x^2+3xy+y^2))=0. \end{gathered}$$

Consequently for $\epsilon=-1$ and $s\in[2,\eta)$ G attains its negative minimum in F on the line $x=-y$. It follows that if $g\leq 1$ then $x=-y\in(0,1), E=0$ and $\epsilon=1$ or $\epsilon=0$. These solutions are also valid for $g\geq 2$ but in this case we have also solutions with $x\neq -y$ both for $\epsilon=-1$ and $s\in(0,\eta)$ (then $-1<y<x<1$) and for $\epsilon=1$ and $s\in(0,2)$ (in this case there exist solutions with $1<y<x$). Now we give explicit formulas for the case $x=-y\in(0,1)$. Then we get

$$\begin{aligned} P(t) &\\ = &\frac{1}{x(15-5x^2-11x^4+x^6)}((t^2-x^2)(s(-15+10x^2-3x^4+t^2(10+12x^2-6x^4)\\ &+t^4(-3-6x^2+x^4))+4\epsilon x(x^2(-5+x^2)-t^4(3+x^2)+t^2(5+2x^2+x^4)))). \end{aligned} \tag{2.14}$$

Thus

$$P(0)=\frac{-4\epsilon x^4(x^2-5)+sx(15-10x^2+3x^4)}{15-5x^2-11x^4+x^6}$$

and, since $\lim_{x\to 0^+}\frac{P(0)}{x}=s>0$, there exists $\epsilon_s>0$ such that $P(t)>0$ if $t\in(0,x)$ for all $x\in(0,\epsilon_s)$. In fact in the case $\epsilon=-1$ the real number ϵ_s is the first positive root of the polynomial $-4x^3(x^2-5)+s(-15+10x^2-3x^4)$ and $\epsilon_s=1$ if $\epsilon=1$. Note also that in both cases $\epsilon_s=1$ if $s\geq 2$. Now the function $z_0(t)=\frac{1}{(1-t^2)}P(t)$ is positive on $(-x,x), x\in(0,\epsilon_s)$. If $x\in(0,\epsilon_s)$ then there exists a solution $h:(-a,a)\to(-sx,sx)$, where

$$a=\lim_{t\to sx^-}\int_0^t\frac{dh}{\sqrt{z_0(\frac{h}{s})}},$$

of an equation

$$h'=\sqrt{z_0(\frac{h}{s})},$$

such that $h(-a)=-sx, h(a)=sx, h'(-a)=h'(a)=0, h''(-a)=1, h''(a)=-1$. It follows that functions $f=h', g=\sqrt{s^2-h^2}$ are smooth on $(-a,a)$ and satisfy the boundary conditions described in Th.2. Consequently the metric

$$g_x = dt^2 + f(t)^2\theta^2 - g(t)^2 p^* g_{can},$$

on the manifold $(-a,a)\times P_k$ extends to the smooth metric on the compact ruled surface $M = P_k \times_{S^1} S^2$ which is a $2-$sphere bundle over Riemannian surface Σ_g. Note that $g(-a)=g(a)=s\sqrt{1-x^2}$. ♢

Theorem 3. *On the surfaces* $M_{0,g}=\mathbb{CP}^1\times\Sigma_g$ *where* $g\geq 2$ *there exists a one-parameter family* $\{g_\alpha : \alpha>1\}$ *of bi-Hermitian* $\mathcal{A}C^\perp$*-metrics. The Ricci tensor* $\rho=\rho_\alpha$ *of* (Σ_0, g_α) *is bi-Hermitian and has two eigenvalues, which are everywhere different.*

Proof. Analogous to [J-3]. Let us write $g'=\sqrt{P(g)}$. Then

$$g^2P''(g) - 2P'(g)g - 4P(g) + 16 + 6Cg^2 = 0. \tag{2.15}$$

Consequently

$$P(g) = \frac{A}{g} + Bg^4 + Cg^2 + 4, \tag{2.16}$$

where $A, B\in\mathbb{R}$ are arbitrary. Now let $D=1$ and let us consider the equations (we are looking for unknown real numbers A, B, C and (x,y) where $0<y<x$)

$$\begin{gathered} P(y)=0, P(x)=0, \\ P'(y)=2, P'(x)=-2. \end{gathered} \tag{2.17}$$

Then $y=\frac{4(\alpha-1)(\alpha^2+3\alpha+1)}{\alpha(2\alpha^2+\alpha+2)}$, $x=\alpha y=\frac{4(\alpha-1)(\alpha^2+3\alpha+1)}{(2\alpha^2+\alpha+2)}$ where $\alpha>1$. Note that $x,y>0$ and $C>0, A,B<0$. Let us consider an equation ($P=P_\alpha, h=h_\alpha$ depend on the parameter $\alpha>1$)

$$\frac{d^2h}{dt^2} = \frac{1}{2}P'(h),\ h'(0)=0,\ h(0)=y=\frac{4(\alpha-1)(\alpha^2+3\alpha+1)}{\alpha(2\alpha^2+\alpha+2)}. \tag{2.18}$$

This equation is equivalent to (if $t\in D=\{t\geq 0 : h'(t)\geq 0\}$)

$$\frac{dh}{dt} = \sqrt{P(h)},\quad h(0)=y=\frac{4(\alpha-1)(\alpha^2+3\alpha+1)}{\alpha(2\alpha^2+\alpha+2)}. \tag{2.19}$$

It follows that $P=P_\alpha$, where $\alpha>1$, has exactly two positive roots $\{x,y\}$ and $P(t)>0$ if $t\in(x,y)$. Note that equation (2.18) admits a smooth periodic solution h defined on the whole of $\mathbb{R}$ and such that im $h=[x,y]$. Now it is easy to check that $\lambda=-10Bh^2-3C$ and $\mu=-5Bh^2-3C$. The tensor $\rho-\frac{\tau}{3}g$ is a Killing tensor with eigenvalues $C, 5Bh^2+C$ corresponding to $\mathcal{D}, \mathcal{D}^\perp$ respectively. Then we obtain a one parameter family of bi-Hermitian $\mathcal{A}C^\perp$-metrics $\{g_\alpha : \alpha>1\}$ on $M_{0,g}$. ♢

3. Einstein neutral bi-Hermitian surfaces. Let us note that the solutions $(M_{k,g}, g_x)$ with $D(x) = 0$ correspond to Einstein neutral surfaces. Let us recall that

$$D = \frac{5(-6s - 24\epsilon x - 12sx^2 - 8\epsilon x^3 + 2sx^4)}{2(-1+x)x(1+x)(15+10x^2-x^4)} \tag{3.1}$$

Consequently there exist Einstein metrics in the family of Gray metrics $(M_{k,g}, g_x)$ if and only if the equation

$$-6s - 24\epsilon x - 12sx^2 - 8\epsilon x^3 + 2sx^4 = 0 \tag{3.2}$$

has a real root $x \in (0,1)$. Now it is not difficult to check that in the case $g = 0$ we have $\epsilon = 1$ and $s \in \mathbb{N}$ and that equation (3.2) does not have any solution in $(0,1)$. Similarly in the case $g = 1$. In the case $g > 1$ we have $s = \frac{k}{g-1} > 0$ and $\epsilon = -1$. In that case equation (3.2) has a real root in $(0,1)$ if and only if $s \in (0,2)$. To show this, let us denote

$$Q(x) = -6s - 24\epsilon x - 12sx^2 - 8\epsilon x^3 + 2sx^4.$$

Then

$$Q''(x) = -24(s + 2\epsilon x - sx^2).$$

Consequently $Q''(x) < 0$ in $(0,1)$ for $\epsilon \geq 0$. In the case where $\epsilon = -1$, since $Q''(0) = -24s < 0$ and $Q''(1) = 48 > 0$, the equation $Q''(x) = 0$ has exactly real one root, say α_s. (In fact, $\alpha_s = s(\sqrt{1+s^2}+1)^{-1}$.) Then, we have $Q'(\alpha_s)(= 16(2-(1+s^2)(\sqrt{1+s^2}+1)^{-1} > 0$ if and only if $(0 <)s < (\sqrt{3}/2)^{\frac{1}{2}}(\sqrt{3}+1)$. In particular, for $0 < s < 2(< (\sqrt{3}/2)^{\frac{1}{2}}(\sqrt{3}+1)$, $Q'(x) \geq Q'(\alpha_s) > 0$, and hence $Q(x)$ is monotone increasing, in $(0,1)$. Since $Q(0) = -6s < 0$ and $Q(1) = 32 - 16s$, it follows that $Q(x)$ has exactly one root in $(0,1)$ if $s \in (0,2)$. On the other hand, if $s/ge2$, then, since $3 + 6x^2 - x^4 > 0$ and $x^4 < x^3$ in $(0,1)$, we see that

$$\begin{aligned} Q(x) &= 24x + 8x^3 - 2s(3 + 6x^2 - x^4) \leq 24x + 8x^3 - 4(3 + 6x^2 - x^4) \\ &< 24x + 8x^3 - 12 - 24x^2 + 4x^3 = 12(x^3 - 2x^2 + 2x - 1) = \\ &12(x^3 - 2x^2 + 2x - 1) = 12(x-1)(x^2 - x + 1) < 0. \end{aligned}$$

Thus the equation $Q(x) = 0$ has exactly one real root $x \in (0,1)$ if and only if $s \in (0,2)$. Since $s = \frac{k}{g-1}$ it follows that the real root $x \in (0,1)$ exists if and only if $k \in \{1, 2, 3, .., 2g-3\}$. Note that for $D = 0$ the polynomial P is even and of degree 4 with exactly one positive root. Consequently we get an Einstein metric on $M_{k,g}$ for all $k \in \{1, 2, .., 2g-3\}$. Thus we obtain on the ruled surfaces $M_{k,g}$ of genus $g > 1$ constructed above exactly $2g-3$ different Einstein neutral bi-Hermitian metrics, exactly one Einstein metric on every surface $M_{k,g}$ for $k \in \{1, 2, ..., 2g-3\}$. Note that for $g = 2$ we obtain only one metric corresponding to the Riemannian Einstein Bergery-Page metric on the first Hirzebruch surface F_1. Consequently we have proved

Theorem 4. *On a ruled surface* $M_{k,g}$ *of genus* $g \geq 2$ *for* $k \in \{1, 2, ..., 2g-3\}$ *there exists an Einstein bi-Hermitian non-Kähler neutral metric.*

Note that in the Riemannian case we have the Page metric which is only one Einstein co-homogeneity 1 Hermitian, non-Kähler metric. (see [P],[B],[LeB],[S]) .

4. Neutral Kähler Gray surfaces. The solutions with $A = 0$ analogously as in the Riemannian case give neutral Kähler Gray surfaces. We shall prove

Theorem 5. *On a ruled surface* $M_{k,g}$ *of genus* $g \geq 2$ *for* $k \in \{1, 2, ..., 2g - 3\}$ *there exists bi-Hermitian Kähler neutral metric.*

Proof. Since $A = 0$ we get $f = \frac{gg'}{s}$. Consequently from (2.3) we get an equation

$$-2\frac{g''}{g} - 4(\frac{g'}{g})^2 - \frac{K}{g^2} = Dg^2 + C. \tag{4.1}$$

Let us write $g' = \sqrt{P(g)}$. Then we can rewrite (4.1) as

$$P'(g) + \frac{4}{g}P(g) + \frac{K}{g} + Dg^3 + Cg = 0, \tag{4.2}$$

with the boundary conditions

$$\frac{1}{2}yP'(y) = s, \frac{1}{2}xP'(x) = -s, \tag{4.3}$$

where $g(a) = y < g(b) = x$, $P(x) = P(y) = 0$ and $P(t) > 0$ for $t \in (y, x)$.

Consequently $P(g) = -\frac{D}{8}g^4 - \frac{C}{6}g^2 + \frac{E}{g^4} - \frac{K}{4}$. Now exactly as in [J-2] we see that the solution satisfying the boundary conditions exists if $K = -4$ and $(s + 2)y^4 + (s - 2)x^4 = 0$. This condition implies that $s < 2, C = 0$, $D > 0$ and $y = \sqrt[4]{\frac{2(2-s)}{D}}, x = \sqrt[4]{\frac{2(2+s)}{D}}$ and $E = \frac{s^2-4}{2D} < 0$. In fact it is not difficult to check that a positive solution satisfying the boundary condition exists if and only if $s \in (0, 2)$ hence for $k < 2(g - 1)$ and that the metrics corresponding to the same s with different D are homothetic.$\diamondsuit$

Note that the situation in the case of neutral 4-manifolds is again quite different from Riemannian case where we have only one irreducible non-Einstein compact Kähler Gray surface (see [A-C-G],[J-2]).

The author is very grateful to the referee for the valuable comments and pointing out a mistake in the first version of the paper.

References.

[A] V. Apostolov *Generalized Goldberg-Sachs theorems for pseudo-Riemannian four-manifolds* J. Geom. Phys. 27,(1998), 185-198.

[A-C-G] V. Apostolov, D. Calderbank and P. Gauduchon *The geometry of weakly selfdual Kähler surfaces* Compositio Math. 135,(2003),179-322.

[B] Bérard-Bergery, L.*Sur de nouvelles variétés riemanniennes d'Einstein*, Institut Élie Cartan,6, Univ. Nancy, (1982), 1-60.

[Be] A. Besse, *Einstein manifolds*, Springer Verlag, 1987.

[D] M. Dunajski, *Anti-self-dual four manifolds with a parallel real spinor*, Proc. R. Soc. London. A(2002) 458, 1205-1222

[G] A. Gray *Einstein-like manifolds which are not Einstein* Geom. Dedicata **7** (1978) 259-280.

[J-1] W. Jelonek, *Compact Kähler surfaces with harmonic anti-self-dual Weyl tensor*, Diff. Geom. and Appl.16 (2002),267-276

[J-2] W. Jelonek, *Extremal Kähler $\mathcal{AC}^{\perp}$-surfaces*, Bull. Belg. Math. Soc. - Simon Stevin, 9,(2002), 561-571.

[J-3] W. Jelonek, *Bi-Hermitian Gray surfaces* , (to appear in Pacific Math. J)

[J-4] W. Jelonek, *Bi-Hermitian Gray surfaces II*, Preprint 2003

[K] H. Kamada, *Self-dual Kähler metrics of neutral signature on complex surfaces* Tohoku Mathematical Publications, September 2002.

[LeB] C. Lebrun *Einstein metrics on complex surfaces*, in Geometry and Physics (Aarhus 1995) eds. J. Andersen, J. Dupont, H. Pedersen and A. Swann, Lect. notes in Pure Appl. Math., Marcel Dekker, 1996.

[M] Y. Matsushita *Fields of 2-planes and two kinds of almost complex structures on compact 4-dimensional manifolds* Math. Zeitschrift **207** (1991), 281-291. par

[M-L] Y. Matsushita and P. Law *Hitchin-Thorpe Type Inequalities for Pseudo-Riemannian 4-Manifolds of Metric Signature* $(++--)$ Geom. Dedicata **87** (2001), 65-89.

[M-S] B.Madsen, H. Pedersen, Y. Poon, A. Swann *Compact Einstein-Weyl manifolds with large symmetry group.* Duke Math. J. **88** (1997), 407-434.

[ON] B. O'Neill *Semi-Riemannian Geometry*, Academic Press, 1983.

[P] D. Page *A compact rotating gravitational instanton*, Phys. Lett.79 **B** (1978), 235-238.

[Pe] D. Petean *Indefinite Kahler-Einstein Metrics on Compact Complex Surfaces*, Comm. Math. Phys. 189 (1997), 227-235.

[S] P. Sentenac ,*Construction d'une métriques d'Einstein sur la somme de deux projectifs complexes de dimension 2*, Géométrie riemannienne en dimension 4 (Séminaire Arthur Besse 1978-1979) Cedic-Fernand Nathan, Paris (1981), pp. 292-307.

Institute of Mathematics
Technical University of Cracow
Warszawska 24
31-155 Krak/ow,POLAND.
E-mail address: wjelon@usk.pk.edu.pl